

03-08-0

JC09 Rec'd PCT/PTO 07 MAR 2001

FORM PTO-1390 (Modified) (REV 10-95)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER BB-1240	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR) 09/786675	
				MAR 07 2001	
INTERNATIONAL APPLICATION NO. PCT/US99/20419		INTERNATIONAL FILING DATE 7 SEPTEMBER 1999 (07.09.99)		PRIORITY DATE CLAIMED 8 SEPTEMBER 1998 (08.09.98)	
TITLE OF INVENTION PLANT FARNESYLTRANSFERASES					
APPLICANT(S) FOR DO/EO/US CAHOON, Rebecca E., et al.					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information					
<ol style="list-style-type: none">1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.3. <input checked="" type="checkbox"/> This is an express request to 'bypass' national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.5. <input checked="" type="checkbox"/> A copy of the International Application was filed (35 U.S.C. 371 (c) (2))<ol style="list-style-type: none">a. <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau.b. <input type="checkbox"/> has been transmitted by the International Bureau.c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US)6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371 (c) (2)).7. <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210).8. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c) (3))<ol style="list-style-type: none">a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).b. <input type="checkbox"/> have been transmitted by the International Bureau.c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.d. <input checked="" type="checkbox"/> have not been made and will not be made.9. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).10. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).11. <input type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409)12. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).					
Items 13 to 18 below concern document(s) or information included :					
<ol style="list-style-type: none">13. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.14. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.15. <input type="checkbox"/> A FIRST preliminary amendment. A SECOND or SUBSEQUENT preliminary amendment.16. <input type="checkbox"/> A substitute specification.17. <input checked="" type="checkbox"/> A change of power of attorney and/or address letter.18. <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail.19. <input type="checkbox"/> Other items or information:					
<div style="border: 1px solid black; padding: 5px;"><p>17. General Power of Attorney</p><p>18. Express Mailing Label No.: EL031052405US</p></div>					

APPLICATION NO. (SEE INSTRUCTIONS) **09/786675**

INTERNATIONAL APPLICATION NO.
PCT/US99/20419

ATTORNEY'S DOCKET NUMBER
BB-1240

20. The following fees are submitted

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :

- ☒ Search Report has been prepared by the EPO or JPO \$860.00
- ☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) \$690.00
- ☐ No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$760.00
- ☐ Neither international preliminary examination fee paid to USPTO (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$1000.00
- ☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) \$100.00
- And all claims satisfied provisions of PCT Article 33(2)-(4)

CALCULATIONS PTO USE ONLY

ENTER APPROPRIATE BASIC FEE AMOUNT =

\$860.00

Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)).

☐ 20 ☐ 30

\$0.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	34 - 20 =	14 x	\$18.00	\$252.00
Independent Claims	6 - 3 =	14 x	\$80.00	\$240.00

Multiple Dependent Claims (check if applicable)

☒

\$270.00

TOTAL OF ABOVE CALCULATIONS =

\$762.00

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable).

☐

\$0.00

SUBTOTAL =

\$762.00

Processing Fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)).

☐ 20 ☐ 30

\$0.00

TOTAL NATIONAL FEE =

\$1,622.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).

☐

\$0.00

TOTAL FEES ENCLOSED =

\$1,622.00

Amount to be :
refunded \$

Charged \$

☐ A check in the amount of _____ to cover the above fees enclosed.

☒ Please charge my Deposit Account No. **04-1928** in the amount of **\$1,622.00** to cover the above fees.

☒ The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. **04-1928** a duplicate copy of this sheet is enclosed.

NOTE : Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (CFR 1.37(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

**RIZZO, Thomas M.
E. I. DU PONT DE NEMOURS AND COMPANY
Legal Patent Records Center
1007 Market Street
Wilmington, Delaware 19898
United States of America**

Andrew L. Schaeffer Reg. No. 33,645
SIGNATURE Andrew L. Schaeffer
for **RIZZO, THOMAS M.**

NAME

41,272

REGISTRATION NUMBER
MARCH 6, 2001

DATE



10 Rec'd PCT/PTC 04 JUN 2001

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the Application of:

REBECCA E. CAHOON

CASE NO.: BB1240

APPLICATION NO.: 09/786,675

GROUP ART UNIT: UNKNOWN

INTERNATIONAL FILING DATE: 07 SEP 1999 EXAMINER: UNKNOWN

FOR: PLANT FARNESYLTRANSFERASES

**PRELIMINARY AMENDMENT AND RESPONSE TO NOTIFICATION OF
MISSING REQUIREMENTS UNDER 35 U.S.C. 371**

Assistant Commissioner for Patents
Box PCT
Washington, D.C. 20231

Sir:

In response to a Notification of Missing Requirements under 35 U.S.C. 371 dated
10 April 2001, please amend the application as follows.

IN THE SPECIFICATION

Please replace the originally filed Sequence Listing with the attached substitute
Sequence Listing.

REMARKS

Please charge any necessary fee to Deposit Account 04-1928 E. I. du Pont de Nemours
and Company).

In view of the foregoing, allowance of the above-referenced application is respectfully
requested.

Respectfully submitted,

Thomas M. Rizzo, Ph.D.
Attorney For Applicants
Registration No. 41,272
Telephone: 302-892-7760
Facsimile: 302-892-1026

Dated: May 30, 2001

Figure 1 consists of seven sub-diagrams labeled 1 through 7, illustrating the steps of the proposed algorithm for finding the minimum value in a 2D array. Each diagram shows a 2D array with the current minimum value highlighted in red and the search area being updated.

- Diagram 1:** The initial 2D array is shown. The minimum value is 1, located at the bottom-left corner.
- Diagram 2:** The search area is updated to the top-right corner. The minimum value is 2, located at the top-right corner.
- Diagram 3:** The search area is updated to the top-left corner. The minimum value is 3, located at the top-left corner.
- Diagram 4:** The search area is updated to the bottom-right corner. The minimum value is 4, located at the bottom-right corner.
- Diagram 5:** The search area is updated to the bottom-left corner. The minimum value is 5, located at the bottom-left corner.
- Diagram 6:** The search area is updated to the top-right corner. The minimum value is 6, located at the top-right corner.
- Diagram 7:** The search area is updated to the top-left corner. The minimum value is 7, located at the top-left corner.

5

10

10

15

20

25

35

- in plants. These genes may be used in plant cells to control cell growth and produce plants with improved water stress tolerance. Accordingly, the availability of nucleic acid sequences encoding all or a portion of farnesyltransferase proteins would facilitate studies to better understand cell growth in plants, provide genetic tools to control cell growth and improve tolerance to drought in mature plants.

SUMMARY OF THE INVENTION

- The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 300 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn farnesyltransferase polypeptide of SEQ ID NO:2, a rice farnesyltransferase polypeptide of SEQ ID NO:4, a soybean farnesyltransferase polypeptide of SEQ ID NO:6, a soybean farnesyltransferase polypeptide of SEQ ID NO:8, a wheat farnesyltransferase polypeptide of SEQ ID NO:10, a corn farnesyltransferase polypeptide of SEQ ID NO:12, a rice farnesyltransferase polypeptide of SEQ ID NO:14, a soybean farnesyltransferase polypeptide of SEQ ID NO:16, and a soybean farnesyltransferase polypeptide of SEQ ID NO:18. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above. It is preferred that the isolated polynucleotides of the claimed invention consists of regions of the isolated polynucleotide selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, and 17 that codes for the polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 and 18. The present invention also relates to an isolated polynucleotide comprising a nucleotide sequences of at least one of 40 (preferably 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17 and the complement of such nucleotide sequences.

The present invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to suitable regulatory sequences.

- The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eucaryotic, such as a yeast or a plant cell, or procaryotic, such as a bacterial cell. The present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

- The present invention relates to a process for producing an isolated host cell comprising a chimeric gene or isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to a farnesyltransferase polypeptide of at least 300 amino acids that has at least 80% homology based on the Clustal method of alignment

compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16, 18.

The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a farnesyltransferase polypeptide in a plant cell, the method comprising the steps of:

constructing an isolated polynucleotide or chimeric gene of the present invention;

introducing the isolated polynucleotide into a plant cell;
measuring the level of farnesyltransferase polypeptide in the plant cell containing the polynucleotide; and

comparing the level of farnesyltransferase polypeptide in the plant cell containing the isolated polynucleotide with the level of farnesyltransferase polypeptide in a plant cell that does not contain the isolated polynucleotide.

The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a farnesyltransferase gene, preferably a plant farnesyltransferase gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 40 (preferably 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17 and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a farnesyltransferase polypeptide amino acid sequence.

The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a farnesyltransferase protein comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying Sequence Listing which form a part of this application.

Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 1
Farnesyltransferase Subunits

Protein	Clone Designation	SEQ ID NO: (Nucleotide)	(Amino Acid)
Farnesyltransferase Alpha Subunit	cen3n.pk0052.a2	1	2
Farnesyltransferase Alpha Subunit	rls6.pk0017.g2	3	4
Farnesyltransferase Alpha Subunit	src1c.pk002.f15	5	6
Farnesyltransferase Alpha Subunit	sgs2c.pk006.n4	7	8
Farnesyltransferase Alpha Subunit	Contig composed of: wdk2c.pk013.d24 wrl.pk0110.b1	9	10
Farnesyltransferase Beta Subunit	p0127.cntbu.18r	11	12
Farnesyltransferase Beta Subunit	rlr24.pk0007.d6	13	14
Farnesyltransferase Beta Subunit	sfl1.pk0086.h10	15	16
Farnesyltransferase Beta Subunit	sgs2c.pk002.g2	17	18

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, a "polynucleotide" is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, or synthetic DNA.

As used herein, "contig" refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping

sequences. Where common or overlapping sequences exist between two or more nucleic acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

As used herein, "substantially similar" refers to nucleic acid fragments wherein
5 changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to
10 mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention
15 encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100%
20 sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more
25 hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule
30 would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 40 (preferably 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9,
35 11, 13, 15, 17 and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a farnesyltransferase polypeptide in a plant cell.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (polynucleotides) encode amino acid sequences that are 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragment encode amino acid sequences that are 85% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are 95% identical to the amino acid sequences reported herein. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively

identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

"Codon degeneracy" refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

"Synthetic nucleic acid fragments" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. "Chemically synthesized", as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene,

5 comprising regulatory and coding sequences that are not found together in nature.

Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature.

“Endogenous gene” refers to a native gene in its natural location in the genome of an
10 organism. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid
15 sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

20 “Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence which can stimulate promoter activity and may be an innate element of
25 the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at
30 different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82.
35 It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The "translation leader sequence" refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The "3' non-coding sequences" refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into polypeptide by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. "Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production

of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

5 "Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and
10 propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a
15 nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*)
20 may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the
25 transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

30 Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

35 Nucleic acid fragments encoding at least a portion of several farnesyltransferase subunits have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the

same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g.,

5 polymerase chain reaction, ligase chain reaction).

For example, genes encoding other farnesyltransferase alpha or beta subunits, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific

10 oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed

15 and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in

20 polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA

25 precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed

30 from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a

35 nucleotide sequence of at least one of 40 (preferably 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17 and the complement of such nucleotide sequences may be used in such

methods to obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a thioredoxin polypeptide.

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of farnesyltransferase activity in those cells.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by altering the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant*

Phys.100:1627-1632) added and/or with targeting sequences that are already present removed. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

- It may also be desirable to reduce or eliminate expression of genes encoding the
- 5 instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences.
- Alternatively, a chimeric gene designed to express antisense RNA for all or part of the
- 10 instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

- Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant
- 15 phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation
- 20 available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

- 25 The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above,
- 30 it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening large numbers of transformants. The nature of these screens will generally be chosen on practical grounds, and is not an inherent part of the invention. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein
- 35 encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell

5 extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate
10 microorganisms via transformation to provide high level expression of the encoded farnesyltransferase subunit. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 7).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part
15 of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then
20 be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted
25 and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous
30 publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

Nucleic acid probes derived from the instant nucleic acid sequences may also be used
35 for physical mapping (i.e., placement of sequences on physical maps; *see* Hoheisel et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 110 Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various corn, rice, soybean and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

15 cDNA Libraries from Corn, Rice, Soybean and Wheat

Library	Tissue	Clone
cen3n	Corn endosperm 20 days after pollination*	cen3n.pk0052.a2
	Corn nucellus tissue, 5 days after silking*	p0127.cntbu18r
rlr24	Rice leaf 15 days after germination, 24 hours after infection of strain <i>Magnaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr24.pk0007.d6
rls6	Rice leaf 15 days after germination, 6 hours after infection of <i>Magnaporthe grisea</i> strain 4360-R-67 (AVR2-YAMO); susceptible	rls6.pk0017.g2
		sfl1.pk0086.h10
sgs2c	Soybean seeds 14 hours after germination	sgs2c.pk006.n4
sgs2c	Soybean seeds 14 hours after germination	sgs2c.pk002.g2
src1c	Soybean 8 day old root infected with cyst nematode <i>Heterodera glycinesis</i>	src1c.pk002.fl5
wdk2c	Wheat developing kernel, 7 days after anthesis	wdk2c.pk013.d24
wr1	Wheat root from 7 day old seedling	wr1.pk0110.b1

*These libraries were normalized essentially as described in U.S. Patent No. 5,482,845, incorporated herein by reference.

20 cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be

introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

EXAMPLE 2

Identification of cDNA Clones

cDNA clones encoding farnesyltransferase subunits were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding Farnesyltransferase Alpha Subunits

The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to farnesyltransferase alpha subunit from *Pisum sativum* (NCBI Identifier No. gi 2246442) and *Arabidopsis thaliana* (NCBI Identifier No. gi 3142698). Shown in Table 3 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), or contigs assembled from two or more ESTs ("Contig"):

TABLE 3

BLAST Results for Sequences Encoding Polypeptides Homologous
to *Pisum sativum* and *Arabidopsis thaliana* Farnesyltransferase Alpha Subunits

Clone	Status	BLAST pLog Score
cen3n.pk0052.a2	FIS	115.00 (gi 2246442)
rls6.pk0017.g2	FIS	121.00 (gi 3142698)
src1c.pk002.f15	FIS	152.00 (gi 2246442)
sgs2c.pk006.n4	FIS	150.00 (gi 2246442)
Contig composed of: wdk2c.pk013.d24 wr1.pk0110.b1	Contig	114.00 (gi 3142698)

- 5 The data in Table 4 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8 and 10 and the *Pisum sativum* and *Arabidopsis thaliana* sequences (SEQ ID NO:19 and 20 respectively). The percent identity between the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8 and 10 ranged between 55-99%.

10

TABLE 4

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences
of cDNA Clones Encoding Polypeptides Homologous
to *Pisum sativum* and *Arabidopsis thaliana* Farnesyltransferase Alpha Subunits

SEQ ID NO.	Percent Identity to
2	60%
4	62%
6	77%
8	76%
10	62%

- 15 Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for
- 20 pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a farnesyltransferase alpha subunit. These sequences
- 25 alpha subunit.

EXAMPLE 4Characterization of cDNA Clones Encoding Farnesyltransferase Beta Subunits

The BLASTX search using the EST sequences from clones listed in Table 5 revealed similarity of the polypeptides encoded by the cDNAs to farnesyltransferase beta subunit from *Lycopersicon esculentum* (NCBI Identifier No. gi 1815668), *Pisum sativum* (NCBI Identifier No. gi 417482) and *Pisum sativum* (NCBI Identifier No. gi 169049). Shown in Table 5 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), or contigs assembled from two or more ESTs ("Contig"):

TABLE 5

BLAST Results for Sequences Encoding Polypeptides Homologous
to *Lycopersicon esculentum* and *Pisum sativum* Farnesyltransferase Beta Subunits

Clone	Status	BLAST pLog Score
p0127.cntbu18r	FIS	149.00 (gi 1815668)
rlr24.pk0007.d6	FIS	110.00 (gi 1815668)
sfl1.pk0086.h10	FIS	>254.00 (gi 417482)
sgs2c.pk002.g2	EST	62.00 (gi 169049)

The data in Table 6 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOS:12, 14, 16 and 18 and the *Lycopersicon esculentum* (gi 1815668), *Pisum sativum* (gi 417482) and *Pisum sativum* (gi 169049) sequences. The percent identity between the amino acid sequences set forth in SEQ ID NOS:12, 14, 16 and 18 ranged between 50-81%.

TABLE 6

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences
of cDNA Clones Encoding Polypeptides Homologous
to *Lycopersicon esculentum* and *Pisum sativum* Farnesyltransferase Beta Subunits

SEQ ID NO.	Percent Identity to
p0127.cntbu18r	55%
rlr24.pk0007.d6	57%
sfl1.pk0086.h10	77%
sgs2c.pk002.g2	62%

Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for

pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a farnesyltransferase beta subunit. These sequences represent the first corn, rice, soybean and wheat sequences encoding a farnesyltransferase beta subunit.

EXAMPLE 5

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-Sall fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptides, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic

proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

- The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

- The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 μ m in diameter) are coated with DNA using the following technique. Ten μ g of plasmid DNAs are added to 50 μ L of a suspension of gold particles (60 mg per mL). Calcium chloride (50 μ L of a 2.5 M solution) and spermidine free base (20 μ L of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 μ L of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 μ L of ethanol. An aliquot (5 μ L) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

- For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

- Seven days after bombardment the tissue can be transferred to N6 medium that contains glufosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-

supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

- Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

EXAMPLE 6

Expression of Chimeric Genes in Dicot Cells

- A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), SmaI, KpnI and XbaI. The entire cassette is flanked by Hind III sites.

- The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

- Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

- Soybean embryogenic suspension cultures can maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent

No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 μ L of a 60 mg/mL 1 μ m gold particle suspension is added (in order): 5 μ L DNA (1 μ g/ μ L), 20 μ L spermidine (0.1 M), and 50 μ L CaCl_2 (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five μ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 7

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter

system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence encoding a first polypeptide of at least 300 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn farnesyltransferase polypeptide of SEQ ID NO:2, a rice farnesyltransferase polypeptide of SEQ ID NO:4, a soybean farnesyltransferase polypeptide of SEQ ID NO:6, a soybean farnesyltransferase polypeptide of SEQ ID NO:8, a wheat farnesyltransferase polypeptide of SEQ ID NO:10, a corn farnesyltransferase polypeptide of SEQ ID NO:12, a rice farnesyltransferase polypeptide of SEQ ID NO:14, a soybean farnesyltransferase polypeptide of SEQ ID NO:16, and a soybean farnesyltransferase polypeptide of SEQ ID NO:18.

2. An isolated polynucleotide comprising the complement of the polynucleotide of Claim 1.

3. The isolated polynucleotide of Claim 1, wherein the nucleotide sequence comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15 and 17 that codes for the polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 and 18.

4. The isolated polynucleotide of Claim 1 which is DNA.

5. The isolated polynucleotide of Claim 1 which is RNA.

6. A chimeric gene comprising the isolated polynucleotide of Claim 1 or Claim 2 operably linked to suitable regulatory sequences.

7. An isolated host cell comprising the chimeric gene of Claim 6.

8. An isolated host cell comprising an isolated polynucleotide of Claim 1.

9. The isolated host cell of Claim 8, wherein the host cell is yeast.

10. The isolated host cell of Claim 8, wherein the host cell is a bacterial cell.

11. The isolated host cell of Claim 8, wherein the host cell is a plant cell.

12. A virus comprising the isolated polynucleotide of Claim 1.

13. A process for producing an isolated host cell comprising the chimeric gene of Claim 6, the process comprising either transforming or transfecting an isolated compatible host cell with the chimeric gene of Claim 6.

14. A farnesyltransferase polypeptide of at least 300 amino acids that has at least 80% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 and 18.

15. A method of selecting an isolated polynucleotide that affects the level of expression of a farnesyltransferase polypeptide in a plant cell, the method comprising the steps of:

constructing an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17 and the complement of such nucleotide sequences;

- 5 introducing the isolated polynucleotide into a plant cell;
- measuring the level of farnesyltransferase polypeptide in the plant cell containing the polynucleotide; and
- comparing the level of farnesyltransferase polypeptide in the plant cell containing the isolated polynucleotide with the level of farnesyltransferase polypeptide in a
- 10 plant cell that does not contain the polynucleotide.

16. The method of Claim 15 wherein the isolated polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, and 17 that codes for the polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 and 18.

- 15 17. The method of Claim 15 wherein the isolated polynucleotide is DNA.
18. The method of Claim 15 wherein the isolated polynucleotide is RNA.
19. The method of Claim 15 wherein the isolated polynucleotide is a chimeric gene comprising the nucleotide sequence operably linked to suitable regulatory sequences.
20. A method of selecting an isolated polynucleotide that affects the level of
- 20 expression of farnesyltransferase polypeptide in a plant cell, the method comprising the steps of:

 constructing the isolated polynucleotide of Claim 1;

 introducing the isolated polynucleotide into a plant cell;

 measuring the level of farnesyltransferase polypeptide in the plant cell

- 25 containing the polynucleotide; and
- comparing the level of farnesyltransferase polypeptide in the plant cell containing the isolated polynucleotide with the level of farnesyltransferase polypeptide in a plant cell that does not contain the isolated polynucleotide.

21. A method of obtaining a nucleic acid fragment encoding a substantial portion of
- 30 a farnesyltransferase gene comprising the steps of:

 synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 40 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17 and the complement of such nucleotide sequences; and

- 35 amplifying a nucleic acid sequence using the oligonucleotide primer.

22. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a farnesyltransferase protein comprising the steps of:

probing a cDNA or genomic library with an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17, and the complement of such nucleotide sequences;

- 5 identifying a DNA clone that hybridizes with the isolated polynucleotide;
 isolating the identified DNA clone; and
 sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

23. The isolated polynucleotide of Claim 1, wherein the first polypeptide is
10 compared to the corn farnesyltransferase polypeptide of SEQ ID NO:2.

24. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the rice farnesyltransferase polypeptide of SEQ ID NO:4.

25. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the soybean farnesyltransferase polypeptide of SEQ ID NO:6.

15 26. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the soybean farnesyltransferase polypeptide of SEQ ID NO:8.

27. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the wheat farnesyltransferase polypeptide of SEQ ID NO:10.

28. The isolated polynucleotide of Claim 1, wherein the first polypeptide is
20 compared to the corn farnesyltransferase polypeptide of SEQ ID NO:12.

29. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the rice farnesyltransferase polypeptide of SEQ ID NO:14.

30. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the soybean farnesyltransferase polypeptide of SEQ ID NO:16.

25 31. The isolated polynucleotide of Claim 1, wherein the first polypeptide is compared to the soybean farnesyltransferase polypeptide of SEQ ID NO:18.

DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

PLANT FARNESYLTRANSFERASES

the specification of which is attached hereto unless the following box is checked:

■ was filed on **07 SEPTEMBER 1999** as U.S. Application No. _____ or PCT International Application No. **PCT/US99/20419** and was amended on _____ (# applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment received to above.

I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Application No.

Country

Filing Date

Priority Claimed (Yes/No)

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States Provisional Application(s) listed below.

U.S. Provisional Application No.

U.S. Filing Date

60/099,521

08 SEPTEMBER 1998

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International Application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application or PCT International Application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application No.

Filing Date

Status (patented, pending or abandoned)

POWER OF ATTORNEY: I hereby appoint the following attorney(s) and/or agent(s) the power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:Name: **WILLIAM R. MAJARIAN**Registration No.: **41,173**

Send correspondence and direct telephone calls to:

WILLIAM R. MAJARIAN
E. I. du Pont de Nemours and Company
Legal - Patents
Wilmington, DE 19898, U.S.A.
Tel. No.
(302) 992-4926
Fax No.
(302) 892-7949

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

INVENTOR(S)

Full Name of Inventor	Last Name CAHOON	First Name REBECCA	Middle Name E.
	Signature (please sign full name): <i>Rebecca E. Cahoon</i>		Date: <i>10/18/99</i>
Residence & Citizenship	City WILMINGTON <i>DE</i>	State or Foreign Country DELAWARE	Country of Citizenship U.S.A.
Post Office Address	Post Office Address 2331 WEST 18TH STREET	City WILMINGTON	State or Country DELAWARE Zip Code 19806
Full Name of Inventor	Last Name MIAO	First Name GUO-HUA	Middle Name
	Signature (please sign full name): <i>Guo-Hua Miao</i>		Date: <i>10/19/99</i>
Residence & Citizenship	City HOCKESSIN <i>DE</i>	State or Foreign Country DELAWARE	Country of Citizenship CN
Post Office Address	Post Office Address 202 CHERRY BLOSSOM PLACE	City HOCKESSIN	State or Country DELAWARE Zip Code 19707
Full Name of Inventor	Last Name POWELL	First Name WAYNE	Middle Name
	Signature (please sign full name): <i>Wayne Powell</i>		Date: <i>10.19.99.</i>
Residence & Citizenship	City WILMINGTON <i>DE</i>	State or Foreign Country DELAWARE	Country of Citizenship GB
Post Office Address	Post Office Address 718 MOUNT LEBANON ROAD	City WILMINGTON	State or Country DELAWARE Zip Code 19803

☐ Additional Inventors are being named on separately numbered sheets attached hereto.

GENERAL POWER OF ATTORNEY
(Concerning Several International Patent Applications)

The undersigned, Vernon R. Rice, Vice President and Assistant General Counsel of E. I. DU PONT DE NEMOURS AND COMPANY, 1007 Market Street, Wilmington, Delaware 19898 USA ("DuPont"), hereby confirms that the power to sign for DuPont has been granted to various individuals (as set forth in the attached excerpt from DuPont's Patent Board Rules of Procedure (January 1988), Appendix Section III.A.4), including the Chairman, Vice-Chairman, and those individuals who are Assistant Secretaries of the Patent Board. Currently these Assistant Secretaries are:

Roger A. Bowman
Linda J. Davis
John E. Griffiths

Miriam D. Meconnahey
Dorothy W. Shafer
Deborah A. Meginniss

In addition, the authority to act on behalf of DuPont before the competent International Authorities in connection with any and all international patent applications filed by it with the United States as Receiving Office and to make or receive payments on its behalf is hereby granted to:

Beardell, Lori Y.	34,293	Katz, Elliott A.	26,396
Belopolsky, Inna	43,319	Kelly, Patricia L.	39,247
Benjamin, Steven C.	36,087	King, Karen K.	34,850
Birch, Linda D.	38,719	Kuller, Mark D.	31,925
Bowen, Jr., Alanson G.	24,027	Kruckel, Charles E.	27,344
Christenbury, Lynne M.	30,971	Jarnholm, Arne R.	30,396
Cotreau, William J.	36,490	Langworthy, John A.	32,255
Deitch, Gerald E.	30,457	Lerman, Bart E.	31,897
Deshmukh, Sudhir	33,677	Levitt, Cary A.	31,848
Dobson, Kevin S.	40,296	Magee, Thomas H.	27,355
Duffy, Roscenne R.	33,869	Mayer, Nancy S.	29,190
Edwards, Mark A.	39,542	Medwick, George M.	27,456
Estrin, Barry	26,452	Morrissey, Bruce W.	30,663
Evans, Craig H.	31,825	Santopietro, Lois A.	36,264
Fair, Tamera L.	33,867	Schaeffer, Andrew L.	33,605
Feltham, S. Neil	36,506	Sebree, Chyrra J.	45,348
Floyd, Linda Axamethy	33,692	Shafer, Robert J.	24,437
Frank, George A.	27,636	Shay, Lucas K.	34,724
Golian, Andrew G.	25,293	Shipley, James E.	32,003
Gorman, Thomas W.	31,959	Siegell, Barbara C.	30,684
Gould, David J.	25,338	Sinnott, Jessica M.	34,015
Griffiths, John E.	32,647	Steinberg, Thomas W.	37,013
Hamby, Jane O.	32,872	Stevenson, Robert B.	26,039
Hamby, William H.	31,521	Strickland, Frederick D.	39,041
Heiser, David E.	31,366	Tessari, Joseph A.	32,177
Hendrickson, John S.	30,847	Tulloch, Rebecca W.	36,297
Jones, Brian C.	37,857	Walker, P. Michael	32,602
Joung, J. Kenneth	41,881	Wang, Chen	38,650

The undersigned ratifies fully all actions already taken by the above-named individuals in accordance with the authority granted hereby.

E. I. DU PONT DE NEMOURS AND COMPANY

By: _____

Vernon R. Rice
Vice President and Assistant General Counsel

Date: _____

SEQUENCE LISTING

<110> Cahoon, Rebecca E.
Miao, Gou-Hau
Powell, Wayne

<120> Plant Farnesyltransferases

<130> BB-1240

<140>

<141>

<150> 60/099,521

<151> September 8, 1998

<160> 23

<170> Microsoft Office 97

<210> 1

<211> 1426

<212> DNA

<213> Zea mays

<400> 1

```
gcacgagaca gcgcaattac ttaagctatt tgtattcggga totgatccaa cctcgttggt 60
cagctgggact catcgcccat ggagcacact aagtcaggcc ccagcagttg gccagaactg 120
gccgacgtgg tgccgggtgcc gcaggacgat gggcctagcc ctgtggtgtc catcgccctat 180
cgagatgact ttcgtgaggt catggattac ttccgcgccc tctacctcac cggtgagcga 240
agccctcgcg ctctccgccc caccgcccag gccatcgagc tcaaccccgg caactacact 300
gtctggcatt tccggcgccc tattctggag tcactagatt ttgatttact agaggagatg 360
aaatttgtcg aaaaaattgc tgaatgcaat ccaaaaaaatt accaaatctg gcaccataag 420
agatgcttgc ctgagaaatt aggcacctggt attgcaaaca aagagcatga attcacaatg 480
aagatacttg ctattgatgc aaaaaattat catgcttggt ctcataggca gtgggttctt 540
caagcgttgg ggggatggga gactgaatta gaatactgtg accacttact taaggagac 600
gtcttcaata attcagcttg gaatcagaga tactttgtta taacaagatc accatttctt 660
gggtggccttg cggcaatgcg tgattcagaa gttagactaca caattgaagc tattctagca 720
aacgctcaga atgaaagccc ctggagggtac ctcaagggtc tatacaaggg tgagaataac 780
ctgctagtag aggacgagcg catctctgct gtttgtttca aggtcctgaa gaatgattgg 840
acttgtgat ttgctttgag tttgctgctc gatcttctct gcactgggtt gcagccttca 900
gatgaactta ggtccactct tgaacaata aggagctccc atccctgaaac cgcggatgat 960
gatcctgcag ccgctgtttg ctgtatccgt cagaaatgtg atccctcgcg ggtaaattat 1020
tggtcttggt tcaaggacac tctttctcag atctcatgac ttcacatggg ttaccocctt 1080
gtccgcgctg gtcggggtc tgtagatag acatgtttta gatagtttca ttggacacc 1140
gaacagagcg gacagagtgt atggctgcta cctctccctg gactgaaagc agtgccttga 1200
acgattttgt ttagtaaaat ttgtgagtgt tactgtccca aacaacacct tatgcaacca 1260
tatttgaaat ttccacatgt aagccttgaat ccagggtgtg ttgttaatgt attacaattg 1320
ccatggggagc ctaaatgaga ccataatca ctccactag agtcggaaga ccgtgtcgag 1380
cagttcactc atatgggtcac ttaagcmeta aaaaaaaaaa aaaaaa 1426
```

<210> 2

<211> 326

<212> PRT

<213> Zea mays

<400> 2

Met Glu His Thr Lys Ser Gly Pro Ser Ser Trp Pro Glu Leu Ala Asp
1 5 10 15

Val Val Pro Val Pro Gln Asp Asp Gly Pro Ser Pro Val Val Ser Ile
 20 25 30
 Ala Tyr Arg Asp Asp Phe Arg Glu Val Met Asp Tyr Phe Arg Ala Leu
 35 40 45
 Tyr Leu Thr Gly Glu Arg Ser Pro Arg Ala Leu Arg Leu Thr Ala Glu
 50 55 60
 Ala Ile Glu Leu Asn Pro Gly Asn Tyr Thr Val Trp His Phe Arg Arg
 65 70 75 80
 Leu Ile Leu Glu Ser Leu Asp Phe Asp Leu Leu Glu Glu Met Lys Phe
 85 90 95
 Val Glu Lys Ile Ala Glu Cys Asn Pro Lys Asn Tyr Gln Ile Trp His
 100 105 110
 His Lys Arg Trp Leu Ala Glu Lys Leu Gly Pro Gly Ile Ala Asn Lys
 115 120 125
 Glu His Glu Phe Thr Met Lys Ile Leu Ala Ile Asp Ala Lys Asn Tyr
 130 135 140
 His Ala Trp Ser His Arg Gln Trp Val Leu Gln Ala Leu Gly Gly Trp
 145 150 155 160
 Glu Thr Glu Leu Glu Tyr Cys Asp His Leu Leu Lys Glu Asp Val Phe
 165 170 175
 Asn Asn Ser Ala Trp Asn Gln Arg Tyr Phe Val Ile Thr Arg Ser Pro
 180 185 190
 Phe Leu Gly Gly Leu Ala Ala Met Arg Asp Ser Glu Val Asp Tyr Thr
 195 200 205
 Ile Glu Ala Ile Leu Ala Asn Ala Gln Asn Glu Ser Pro Trp Arg Tyr
 210 215 220
 Leu Lys Gly Leu Tyr Lys Gly Glu Asn Asn Leu Leu Val Glu Asp Glu
 225 230 235 240
 Arg Ile Ser Ala Val Cys Phe Lys Val Leu Lys Asn Asp Trp Thr Cys
 245 250 255
 Val Phe Ala Leu Ser Leu Leu Leu Asp Leu Leu Cys Thr Gly Leu Gln
 260 265 270
 Pro Ser Asp Glu Leu Arg Ser Thr Leu Glu Thr Ile Arg Ser Ser His
 275 280 285
 Pro Glu Thr Ala Asp Asp Asp Pro Ala Ala Ala Val Cys Cys Ile Leu
 290 295 300
 Gln Lys Cys Asp Pro Leu Arg Val Asn Tyr Trp Ser Trp Phe Lys Asp
 305 310 315 320
 Thr Leu Ser Gln Ile Ser
 325

<210> 3
 <211> 1218
 <212> DNA
 <213> Oryza sativa

<400> 3
 gcacgaggtt ctaacgccgc gcgcgccgc gcgctctccg cagaatctga tcgatggcgc 60
 cgctcgtcac gtcgtcggag ggtgcctccg acgagtgttt gccaccacgc cggcgccgcg 120
 agctggcgga cgtggtcccc gtgacgcagg acgacgggoc ccaccocgtg gtggccatcg 180
 cctaccggga cgaagtccgc gaggtcatgg actacttccg cgcctctcac ttccgcggcg 240
 agcgcagcgt ccgcgccttc cactcacccg ccgaggtcat cgacctaat cccggcaact 300
 acacggtgtg gcattttagg cgtcttgttc tagaggcact ggatgctgat ctgcgtgagg 360
 aaatggattt tgtggaccga attgctgaat gtaacccaaa aaattatcaa atctggcatc 420
 acaagagatg cgttgcggag aaattaggac cagatatgtc aaataaagag cacgaattta 480
 caaggaagat actttctatg gatgctaaaa attaccatgc ttggtctcat aggcagtgagg 540
 ttcttcaagc actgggtgga tgggagactg aactacagta ttgcaaccag ctgcttgagg 600
 aagacgtctt caataattca gcttggaaac agagatacct tgtaataaca agttcaccac 660
 ttcttggagg ccttcgagca atgcgtgact cggaaagtga ttacacagtt ggggctattc 720
 tggctaacc ctcagaatgaa agcccttgga gatacctcaa aggcctgtac aagggtgaaa 780
 ataacttgct gatggctgat gaggcgtatc ctgatgtttg tctcaaggtc ctgaaacatg 840
 attcgacctg cgtatttgct ttgagcttgc tgctcgatct tcttcaaaat gggttataac 900
 cttcagatga actcaaagga actatcgaag caataaagaa cctcgatcct gaagcagatg 960
 aagcagtaga tgcgtatctt gcgactgcaa tctgtcaat attgcagaga tgtgatcccc 1020
 tgccgataaa ttactggctc tgggtacagga ccactatttc ttctcaaaac tgaagcatgc 1080
 agtggcctcc atgaggtcat aatggagata tcttctatct tcgtgtgatt ctgggcgttg 1140
 aggtgcctag ctacatttgt tatgaaatct ccttgggcat aactgatcac tgatattact 1200
 ccaatattgt gttctaaa 1218

<210> 4
 <211> 339
 <212> PRT
 <213> Oryza sativa

<400> 4
 Met Ala Pro Ser Ser Thr Ser Ser Glu Gly Ala Ser Asp Glu Trp Leu
 1 5 10 15
 Pro Pro Ser Arg Arg Pro Glu Leu Ala Asp Val Val Pro Val Thr Gln
 20 25 30
 Asp Asp Gly Pro His Pro Val Val Ala Ile Ala Tyr Arg Asp Glu Phe
 35 40 45
 Arg Glu Val Met Asp Tyr Phe Arg Ala Leu Tyr Phe Ala Gly Glu Arg
 50 55 60
 Ser Val Arg Ala Leu His Leu Thr Ala Glu Val Ile Asp Leu Asn Pro
 65 70 75 80
 Gly Asn Tyr Thr Val Trp His Phe Arg Arg Leu Val Leu Glu Ala Leu
 85 90 95
 Asp Ala Asp Leu Arg Glu Glu Met Asp Phe Val Asp Arg Ile Ala Glu
 100 105 110
 Cys Asn Pro Lys Asn Tyr Gln Ile Trp His His Lys Arg Trp Leu Ala
 115 120 125
 Glu Lys Leu Gly Pro Asp Ile Ala Asn Lys Glu His Glu Phe Thr Arg
 130 135 140

Lys Ile Leu Ser Met Asp Ala Lys Asn Tyr His Ala Trp Ser His Arg
145 150 155 160

Gln Trp Val Leu Gln Ala Leu Gly Gly Trp Glu Thr Glu Leu Gln Tyr
165 170 175

Cys Asn Gln Leu Leu Glu Glu Asp Val Phe Asn Asn Ser Ala Trp Asn
180 185 190

Gln Arg Tyr Leu Val Ile Thr Ser Ser Pro Leu Leu Gly Gly Leu Ala
195 200 205

Ala Met Arg Asp Ser Glu Val Asp Tyr Thr Val Gly Ala Ile Leu Ala
210 215 220

Asn Pro Gln Asn Glu Ser Pro Trp Arg Tyr Leu Lys Gly Leu Tyr Lys
225 230 235 240

Gly Glu Asn Asn Leu Leu Met Ala Asp Glu Arg Ile Ser Asp Val Cys
245 250 255

Leu Lys Val Leu Lys His Asp Ser Thr Cys Val Phe Ala Leu Ser Leu
260 265 270

Leu Leu Asp Leu Leu Gln Ile Gly Leu Gln Pro Ser Asp Glu Leu Lys
275 280 285

Gly Thr Ile Glu Ala Ile Lys Asn Ser Asp Pro Glu Ala Asp Glu Ala
290 295 300

Val Asp Ala Asp Leu Ala Thr Ala Ile Cys Ser Ile Leu Gln Arg Cys
305 310 315 320

Asp Pro Leu Arg Ile Asn Tyr Trp Ser Trp Tyr Arg Thr Thr Ile Ser
325 330 335

Ser Gln Thr

<210> 5

<211> 1261

<212> DNA

<213> Glycine max

<400> 5

gcacgaggat taacgaagga tggaaatctgg gtctagcgaa ggagaagaggy tgcagcaacg 60
cgtgccgttg agggagagag tggagtggtc agatgttact ccggttcctc aaaacgacgg 120
ccctaaccct gtctgtccga tccagtacac tgaagagttt tccgaagtta tggattactt 180
tcgcgccgtt tacctcaccg atgaacgctc ccctcgcgcc ctgctctca cagccgaagc 240
cggtcaattc aactcggca actacactgt gtggcatttc cgacggttgt tacttgagtc 300
gctaaaagtc gacttgaacg atgaactgga tttgtggag cgtatggccg ctggaaaattc 360
taaaaattat cagatgtggc atcatagacg atgggttgcc gagaagttag gtccctgaagc 420
tagaaacaat gagctcgagt tcaccaaaaa gatactgtcc gttgatgcc aacattatca 480
tgcatggctc catagacagt gggctcttca aacactagga ggtatgggaag atgaacttaa 540
ttattgcaca gaactactta aagaagacat ttttaacaat tctgcttga atcagagata 600
ttttgtcata acaaggtctc ctttcttggg gggcctaata gctatgagag agtctgaagt 660
gctttacacc attgaagcca ttatagccta ccctgaaaat gaaagctcgt ggagatatct 720
acgaggactt tataaagggt aaactacttc atgggtaaat gatcctcaag tttcttcagt 780
atgcttaaag attttgagaa ctaagagcaa ctacgtgttt gctcttagca ctattttaga 840

tcttatatgc ttgtgtatc aacaaatga agacattaga gatgccattg acgccttaaa 900
 gaccgcagat atggataaac aagatttaga tgatgatgag aaaggggaac aacaaaattt 960
 aaatatagca cgaatatatt gtctctatcct aaaacaagtt gatccaatta gaaccaacta 1020
 ttggatttgg cgcaagagca gacttcctct atcagcttag taaccaagtt aattaaaggg 1080
 caactctgtg ttatgtgtaa cctagtttat tgaactgga tttttattta ttattatttt 1140
 ttatgtgtgc atgtatctgt ttgtgcaaatt ttatcttttt gtcatgccat tactggcatt 1200
 tgagtgtgaag gattgaaagc catgcagaat aagaaattta agtttttttt tccgttgaaa 1260
 a

<210> 6
 <211> 346
 <212> PRT
 <213> Glycine max

<400> 6
 Met Glu Ser Gly Ser Glu Gly Glu Val Gln Gln Arg Val Pro
 1 5 10 15
 Leu Arg Glu Arg Val Glu Trp Ser Asp Val Thr Pro Val Pro Gln Asn
 20 25 30
 Asp Gly Pro Asn Pro Val Val Pro Ile Gln Tyr Thr Glu Glu Phe Ser
 35 40 45
 Glu Val Met Asp Tyr Phe Arg Ala Val Tyr Leu Thr Asp Glu Arg Ser
 50 55 60
 Pro Arg Ala Leu Ala Leu Thr Ala Glu Ala Val Gln Phe Asn Ser Gly
 65 70 75 80
 Asn Tyr Thr Val Trp His Phe Arg Arg Leu Leu Leu Glu Ser Leu Lys
 85 90 95
 Val Asp Leu Asn Asp Glu Leu Asp Phe Val Glu Arg Met Ala Ala Gly
 100 105 110
 Asn Ser Lys Asn Tyr Gln Met Trp His His Arg Arg Trp Val Ala Glu
 115 120 125
 Lys Leu Gly Pro Glu Ala Arg Asn Asn Glu Leu Glu Phe Thr Lys Lys
 130 135 140
 Ile Leu Ser Val Asp Ala Lys His Tyr His Ala Trp Ser His Arg Gln
 145 150 155 160
 Trp Ala Leu Gln Thr Leu Gly Gly Trp Glu Asp Glu Leu Asn Tyr Cys
 165 170 175
 Thr Glu Leu Leu Lys Glu Asp Ile Phe Asn Asn Ser Ala Trp Asn Gln
 180 185 190
 Arg Tyr Phe Val Ile Thr Arg Ser Pro Phe Leu Gly Gly Leu Lys Ala
 195 200 205
 Met Arg Glu Ser Glu Val Leu Tyr Thr Ile Glu Ala Ile Ile Ala Tyr
 210 215 220
 Pro Glu Asn Glu Ser Ser Trp Arg Tyr Leu Arg Gly Leu Tyr Lys Gly
 225 230 235 240

Glu Thr Thr Ser Trp Val Asn Asp Pro Gln Val Ser Ser Val Cys Leu
245 250 255

Lys Ile Leu Arg Thr Lys Ser Asn Tyr Val Phe Ala Leu Ser Thr Ile
260 265 270

Leu Asp Leu Ile Cys Phe Gly Tyr Gln Pro Asn Glu Asp Ile Arg Asp
275 280 285

Ala Ile Asp Ala Leu Lys Thr Ala Asp Met Asp Lys Gln Asp Leu Asp
290 295 300

Asp Asp Glu Lys Gly Glu Gln Gln Asn Leu Asn Ile Ala Arg Asn Ile
305 310 315 320

Cys Ser Ile Leu Lys Gln Val Asp Pro Ile Arg Thr Asn Tyr Trp Ile
325 330 335

Trp Arg Lys Ser Arg Leu Pro Leu Ser Ala
340 345

<210> 7
<211> 1333
<212> DNA
<213> Glycine max

<400> 7
gcacgaqctt gcgtgtggag tgaagaagat taacgaagga tggaaatctgg gtctagcgaa 60
ggagaagagg tgacagcaacg cgtgccgttg agggagagag tggagtgggtc agatgttact 120
ccggttcctc aaaacgacgg ccctaaccct gtcgttccga tccagtagac tgaagagttt 180
tccgaagtta tggattactt tcgcgcgctt tacctcaccg atgaacgctc cctcgcgcgc 240
ctcgctctca cagccgaagc gtttcaattc aactccggca actacactgt gtggcatttc 300
cgacggttgt tacttgagtc gctaaaagtc gacttgaacg atgaactgga gtttgtggag 360
cgtatggccg ctggaaattc taaaaattat cagatgtggt gtgatgctct gctctgctct 420
ttcttcata ctttgatca tagacgatgg gttgccgaga agttaggtcc tgaagctaga 480
aacaatgagc tcgagttcac caaaaagata ctgtccgttg atgccaaaca ttatcatgca 540
tggtctcata gacagtgggc tcttcaaaca ctaggaggat gggaagatga acttaattat 600
tgacacagaa tacttaaaaga agacattttt aacaattctg cttggaatca gagatatttt 660
gtcataacaa ggtctccttt cttggggggc ctaaaagcta tgagagagtc tgaagtgcct 720
tacaccattg aagccattat agcctaccct gaaaatgaaa gctcgtggag atatctacga 780
ggactttata aaggtgaaac tacttcacgg gtaaatgac ctaagtttc ttcagtatgc 840
ttaaagattt tgagaactaa gagcaactac gtgtttgctc ttgacctatt tttagacttt 900
atatgctttg gttatcaacc aaatgaagac attagagatg ccattgacgc cttaaagacc 960
cgagataggg ataacaaga tttagatgat gatgagaaag gggaaacaaca aaatttaaat 1020
atagcacgaa atatttgctc tatcctaata caagttgatc caattagaac caactattgg 1080
atttggcgca agagcagact tctctatca gcttagtaac caaagtaatt aaagggcaac 1140
cttgtgttat gtgtaaccta gtttattgaa actggatggt tatttattat tattttttat 1200
gttgctatgt atctgtttgt gcaaatattat ctttttgcga tgcattactt ggcatttgag 1260
tgtaaggatt gaaagccatg cagaataaga aatttaagtt tttttttccg ttgaaaaaaa 1320
aaaaaaaaaaa aaa 1333

<210> 8
<211> 358
<212> PRT
<213> Glycine max

<400> 8
Met Glu Ser Gly Ser Ser Glu Gly Glu Glu Val Gln Gln Arg Val Pro
1 5 10 15

Leu Arg Glu Arg Val Glu Trp Ser Asp Val Thr Pro Val Pro Gln Asn
 20 25 30
 Asp Gly Pro Asn Pro Val Val Pro Ile Gln Tyr Thr Glu Glu Phe Ser
 35 40 45
 Glu Val Met Asp Tyr Phe Arg Ala Val Tyr Leu Thr Asp Glu Arg Ser
 50 55 60
 Pro Arg Ala Leu Ala Leu Thr Ala Glu Ala Val Gln Phe Asn Ser Gly
 65 70 75 80
 Asn Tyr Thr Val Trp His Phe Arg Arg Leu Leu Leu Glu Ser Leu Lys
 85 90 95
 Val Asp Leu Asn Asp Glu Leu Glu Phe Val Glu Arg Met Ala Ala Gly
 100 105 110
 Asn Ser Lys Asn Tyr Gln Met Trp Cys Asp Ala Leu Leu Cys Ser Phe
 115 120 125
 Phe His Thr Leu His His Arg Arg Trp Val Ala Glu Lys Leu Gly Pro
 130 135 140
 Glu Ala Arg Asn Asn Glu Leu Glu Phe Thr Lys Lys Ile Leu Ser Val
 145 150 155 160
 Asp Ala Lys His Tyr His Ala Trp Ser His Arg Gln Trp Ala Leu Gln
 165 170 175
 Thr Leu Gly Gly Trp Glu Asp Glu Leu Asn Tyr Cys Thr Glu Leu Leu
 180 185 190
 Lys Glu Asp Ile Phe Asn Asn Ser Ala Trp Asn Gln Arg Tyr Phe Val
 195 200 205
 Ile Thr Arg Ser Pro Phe Leu Gly Gly Leu Lys Ala Met Arg Glu Ser
 210 215 220
 Glu Val Leu Tyr Thr Ile Glu Ala Ile Ile Ala Tyr Pro Glu Asn Glu
 225 230 235 240
 Ser Ser Trp Arg Tyr Leu Arg Gly Leu Tyr Lys Gly Glu Thr Thr Ser
 245 250 255
 Trp Val Asn Asp Pro Gln Val Ser Ser Val Cys Leu Lys Ile Leu Arg
 260 265 270
 Thr Lys Ser Asn Tyr Val Phe Ala Leu Ser Thr Ile Leu Asp Leu Ile
 275 280 285
 Cys Phe Gly Tyr Gln Pro Asn Glu Asp Ile Arg Asp Ala Ile Asp Ala
 290 295 300
 Leu Lys Thr Ala Asp Met Asp Lys Gln Asp Leu Asp Asp Asp Glu Lys
 305 310 315 320
 Gly Glu Gln Gln Asn Leu Asn Ile Ala Arg Asn Ile Cys Ser Ile Leu
 325 330 335

Lys Gln Val Asp Pro Ile Arg Thr Asn Tyr Trp Ile Trp Arg Lys Ser
340 345 350

Arg Leu Pro Leu Ser Ala
355

<210> 9
<211> 1339
<212> DNA
<213> Triticum aestivum

<400> 9
cggaactggc gccgctgccc caggccgacg ggccctgccc cgtcgtctcc atcgcttacc 60
gcggcgactt ccgcgaggtc atggactact tccgcgccct ctacgcgcgc ggcgagcgca 120
gccccgcgcg cctccgcctc accgccgacg coactccacct caacccccgc aactacactg 180
tatggcattt caggcgcggt gttctagagg cactggatgc tgatttattg ctagaaatgc 240
attttgtgga ccaaatgtgt gaattctaac caaaaaatta ccaagtctgg catcacaaaga 300
gatggcttgc tgagaaaata ggaccagatg ctgcaaatag tgaacatgac ttcacaagga 360
agatacttgc tatggatgct aaaaactacc atgcttggtc ccataggcag tgggttcttc 420
aagcattggg tggatgggag agtgaactgc agtactgcaa ccagcttctt gaggaagatg 480
tcttcaataa ctcagcttgg aatcagagat acctgtgtgt aacacgatca ccaattcttg 540
ggggccttgc ggcaatgcgc gactcagaag tagattacac agttgaggcc attatggtga 600
accctcagaa tgaaagcccc tggagatacc tcagagggtt atataaggat gataacaatt 660
tgctgggtggc tgataatgac atttctgatg cttgcctcaa ggtcctgaat aaggattgga 720
catgcgttatt tgctttgagc ttcctgcttg atcttcttcg catgggtttg cagccttcga 780
atgaacttaa aggaaccatc gaagcaatgg agaactctga tcttgaaaac ggacatgctg 840
atatgcagat agctgtctgc tcaatcctgc agaaatgtga tccccgcgcg ataaactact 900
ggatcatggt cagaccactt ctttcttctt agacatctga aaattcagct gaagacagtt 960
ttagcagcat gatgtaactt caatcgaagg ggttgacgca ggtgatgaaa aaccttccct 1020
gtgatcttgg tgccgagcaa tttgtactga ttttactggg aaaaatcaat caatgacagc 1080
atgcccaaca atgtcttctg tgaatatgtt actgcctgat attcacatgt tagcagaatg 1140
agaataacca atcaaaactcc aacgagcaga ttgttacagt aacgcgcact ggtggtgtga 1200
aaatcctgaa atctgcttca gtcactttgc cttgtttaca gttgagctgt ttgttgtgat 1260
ctgtacctaa tgcattgtaca caatcatcaa attattagtt tttgtaccaa tgagtattcg 1320
atgaaaaaaaa aaaaaaaaaa 1339

<210> 10
<211> 309
<212> PRT
<213> Triticum aestivum

<400> 10
Asp Val Ala Pro Leu Pro Gln Ala Asp Gly Pro Cys Pro Val Val Ser
1 5 10 15

Ile Ala Tyr Arg Gly Asp Phe Arg Glu Val Met Asp Tyr Phe Arg Ala
20 25 30

Leu Tyr Ala Ala Gly Glu Arg Ser Pro Arg Ala Leu Arg Leu Thr Ala
35 40 45

Asp Ala Ile His Leu Asn Pro Gly Asn Tyr Thr Val Trp His Phe Arg
50 55 60

Arg Val Val Leu Glu Ala Leu Asp Ala Asp Leu Leu Leu Glu Met His
65 70 75 80

Phe Val Asp Gln Ile Ala Glu Ser Asn Pro Lys Asn Tyr Gln Val Trp
85 90 95

His His Lys Arg Trp Leu Ala Glu Lys Ile Gly Pro Asp Ala Ala Asn
 100 105 110
 Ser Glu His Asp Phe Thr Arg Lys Ile Leu Ala Met Asp Ala Lys Asn
 115 120 125
 Tyr His Ala Trp Ser His Arg Gln Trp Val Leu Gln Ala Leu Gly Gly
 130 135 140
 Trp Glu Ser Glu Leu Gln Tyr Cys Asn Gln Leu Leu Glu Glu Asp Val
 145 150 155 160
 Phe Asn Asn Ser Ala Trp Asn Gln Arg Tyr Leu Val Val Thr Arg Ser
 165 170 175
 Pro Ile Leu Gly Gly Leu Ala Ala Met Arg Asp Ser Glu Val Asp Tyr
 180 185 190
 Thr Val Glu Ala Ile Met Val Asn Pro Gln Asn Glu Ser Pro Trp Arg
 195 200 205
 Tyr Leu Arg Gly Leu Tyr Lys Asp Asp Asn Asn Leu Leu Val Ala Asp
 210 215 220
 Asn Arg Ile Ser Asp Ala Cys Leu Lys Val Leu Asn Lys Asp Trp Thr
 225 230 235 240
 Cys Val Phe Ala Leu Ser Phe Leu Leu Asp Leu Leu Arg Met Gly Leu
 245 250 255
 Gln Pro Ser Asn Glu Leu Lys Gly Thr Ile Glu Ala Met Glu Asn Ser
 260 265 270
 Asp Pro Glu Thr Gly His Ala Asp Ile Ala Val Ala Val Cys Ser Ile
 275 280 285
 Leu Gln Lys Cys Asp Pro Leu Arg Ile Asn Tyr Trp Ser Trp Tyr Gln
 290 295 300
 Thr Thr Leu Ser Ser
 305

<210> 11
 <211> 1359
 <212> DNA
 <213> Zea mays

<400> 11
 atggaccct ccccgagtc gacgccgcc accggagacg acccggcagc ggcggcggat 60
 cccgacctac cgaggctcac ggtgacgcag gtggagcaga tgaagtgga gccagggtt 120
 ggcgacatct accgctccct cttcggggcc gcgccaaca cgaatccat catgctagag 180
 ctgtggcgtg atcagcatat cgagtatctg acgcctgggc tgaggcatat gggaccagcc 240
 tttcatgttc tagatgccaa tcgcccttgg ctatgctact ggatggttca tccacttgct 300
 ttgctggatg aagcacttga tgatgatctt gagaatgata tcatagactt cttagctcga 360
 tgctcaggata aagatgggtg atatatgtgt ggacctggac agttgcctca cctagctacg 420
 acttatgctg ctgtaaatcac acttgtagaca atagggagcg aaagagcatt gtcataatc 480
 aataggggca acctgtacaa ttttatgctg cagatgaaag atgtatcagg tgctttcaga 540
 atgcatgatg gtggcgaaat tgatgtccgt gcttcctaca ccgctatat ggttgccagc 600
 cttgtgaata ttcttgattt taaactggca aaaggtgtag gcgactacat agcaagatgt 660
 caaaccttatg aaggtggtat tgctggggag ccttatgctg aagcacatgg tgggtataca 720

```

ttctgtggat tggctgcttt gatcctgctt aatgaggcag agaaagtga cttgcctagt 780
ttgattggct ggggtgcttt tcgtcaagga gtggaatgcg gatttcaagg acgaactaat 840
aaattggttg atggttgcta ctcttttgg cagggagctg ccattgcttt cacacaaaag 900
ttaattacga ttgttgataa gcaattgaag tcctcgatt cctgcaaaag gccatcagga 960
gaggatgcct gcagcaccag ttcatatggg tgcaccgcga aaaagtcttc ctctgctgtg 1020
gactatgcga agtttggatt tgattttata caacagagca accaaattgg cccactcttc 1080
cataacattg ccctgcaaca atacatecta ctttgtcttc aggtactaga gggaggcttg 1140
agggataaag ctggaaagaa cagagatcac tatattcat gctactgcct cagtggcctc 1200
gcagttagcc agtacagtgc catgactgat actggttcgt gccattacc tcagcatgtg 1260
cttgaccgt actctaattt gctggagcca atccatccac tctacaatgt tgtcctagat 1320
aagtaccata cagcctatga gttcttctca gaagagtga 1359

```

<210> 12

<211> 452

<212> PRT

<213> Zea mays

<400> 12

```

Met Asp Pro Ser Pro Gln Ser Thr Pro Pro Thr Gly Asp Asp Pro Ala
1          5          10          15

```

```

Ala Ala Ala Asp Pro Asp Leu Pro Arg Leu Thr Val Thr Gln Val Glu
20          25          30

```

```

Gln Met Lys Val Glu Ala Arg Val Gly Asp Ile Tyr Arg Ser Leu Phe
35          40          45

```

```

Gly Ala Ala Pro Asn Thr Lys Ser Ile Met Leu Glu Leu Trp Arg Asp
50          55          60

```

```

Gln His Ile Glu Tyr Leu Thr Pro Gly Leu Arg His Met Gly Pro Ala
65          70          75          80

```

```

Phe His Val Leu Asp Ala Asn Arg Pro Trp Leu Cys Tyr Trp Met Val
85          90          95

```

```

His Pro Leu Ala Leu Leu Asp Glu Ala Leu Asp Asp Asp Leu Glu Asn
100         105         110

```

```

Asp Ile Ile Asp Phe Leu Ala Arg Cys Gln Asp Lys Asp Gly Gly Tyr
115         120         125

```

```

Ser Gly Gly Pro Gly Gln Leu Pro His Leu Ala Thr Thr Tyr Ala Ala
130         135         140

```

```

Val Asn Thr Leu Val Thr Ile Gly Ser Glu Arg Ala Leu Ser Ser Ile
145         150         155         160

```

```

Asn Arg Gly Asn Leu Tyr Asn Phe Met Leu Gln Met Lys Asp Val Ser
165         170         175

```

```

Gly Ala Phe Arg Met His Asp Gly Gly Glu Ile Asp Val Arg Ala Ser
180         185         190

```

```

Tyr Thr Ala Ile Ser Val Ala Ser Leu Val Asn Ile Leu Asp Phe Lys
195         200         205

```

```

Leu Ala Lys Gly Val Gly Asp Tyr Ile Ala Arg Cys Gln Thr Tyr Glu
210         215         220

```

Gly Gly Ile Ala Gly Glu Pro Tyr Ala Glu Ala His Gly Gly Tyr Thr
 225 230 235 240
 Phe Cys Gly Leu Ala Ala Leu Ile Leu Leu Asn Glu Ala Glu Lys Val
 245 250 255
 Asp Leu Pro Ser Leu Ile Gly Trp Val Ala Phe Arg Gln Gly Val Glu
 260 265 270
 Cys Gly Phe Gln Gly Arg Thr Asn Lys Leu Val Asp Gly Cys Tyr Ser
 275 280 285
 Phe Trp Gln Gly Ala Ala Ile Ala Phe Thr Gln Lys Leu Ile Thr Ile
 290 295 300
 Val Asp Lys Gln Leu Lys Ser Ser Tyr Ser Cys Lys Arg Pro Ser Gly
 305 310 315 320
 Glu Asp Ala Cys Ser Thr Ser Ser Tyr Gly Cys Thr Ala Lys Lys Ser
 325 330 335
 Ser Ser Ala Val Asp Tyr Ala Lys Phe Gly Phe Asp Phe Ile Gln Gln
 340 345 350
 Ser Asn Gln Ile Gly Pro Leu Phe His Asn Ile Ala Leu Gln Gln Tyr
 355 360 365
 Ile Leu Leu Cys Ser Gln Val Leu Glu Gly Gly Leu Arg Asp Lys Pro
 370 375 380
 Gly Lys Asn Arg Asp His Tyr His Ser Cys Tyr Cys Leu Ser Gly Leu
 385 390 395 400
 Ala Val Ser Gln Tyr Ser Ala Met Thr Asp Thr Gly Ser Cys Pro Leu
 405 410 415
 Pro Gln His Val Leu Gly Pro Tyr Ser Asn Leu Leu Glu Pro Ile His
 420 425 430
 Pro Leu Tyr Asn Val Val Leu Asp Lys Tyr His Thr Ala Tyr Glu Phe
 435 440 445
 Phe Ser Glu Glu
 450
 <210> 13
 <211> 1031
 <212> DNA
 <213> *Oryza sativa*
 <400> 13
 gcacgagggc gtagccgcct ttccgtgaga tcccgcgggc tgcagcgagc tcgcaggccg 60
 cgcccttcgc cgccgcgcac caccgcgcgc atggaccgcc cctcgccgcc gccgcgcgcg 120
 ccatactctc ctgctgctgc tgaggcggt cggcagcggc atagccaggc cgtgagctg 180
 ccccggtgta ctgtgacgca ggtggagcag atgaagtgag aggcgaaggt ggccgaaatc 240
 taccgcgtcc tcttcggcaa cgccccaac gccaatcccc tcatgttaga gctgtggcgt 300
 gaggcagcatg ttgagtattt gacgagaggg ctgaaacatc ttggaccaag ctccatgtg 360
 ctcgatgcc aatcgacctg gctgtgctac tggattatc atgcacttgc tctgttggat 420
 gaaatacctg acgatgttga gcatgatatt gtggacttct tatctcgatg tcaggacaaa 480
 gatggtgtgt atggcggagg acctggacag ttgcctcacc tcgtacaac ttatgtctgt 540

gtaaatcac ttgtaactat agggagtgaa agggcactat catcggtaaa cagggacaac 600
 ctgtacaagt tcatgcttcg gatgaaagat acatcgggag ctttcagaat gcatgatggg 660
 ggtgaaatag atgttcgtgc tagctatact gcaatatcgg ttgccagcct tgtgaacatt 720
 cttgatggg aactagcaaa aggtgttgga aattacataa caagggtgtca aacctatgaa 780
 ggtggcattg ctggggaacc gtatgctgaa gctcatggtg ggtacacttt ttgtgggctg 840
 gctacgatga tcctgcttaa cgaagtggac aaacttgatt tggctagctt gattggctgg 900
 gtggcatttc gccaaaggag ggaatgtgga ttccaaggac gaactaataa attggttgat 960
 ggttgctact ctttttgcca gggagctgct cttgctttaa ctgttcaccg cgtgggcgcc 1020
 actgccaaac g 1031

<210> 14
 <211> 313
 <212> PRT
 <213> Oryza sativa

<400> 14
 Met Asp Pro Pro Ser Pro Pro Pro Pro Pro Pro Tyr Pro Pro Ala Ala
 1 5 10 15
 Ala Glu Gly Gly Pro Ala Ala Asp Ser Gln Ala Ala Glu Leu Pro Arg
 20 25 30
 Leu Thr Val Thr Gln Val Glu Gln Met Lys Val Glu Ala Lys Val Gly
 35 40 45
 Glu Ile Tyr Arg Val Leu Phe Gly Asn Ala Pro Asn Ala Asn Ser Leu
 50 55 60
 Met Leu Glu Leu Trp Arg Glu Gln His Val Glu Tyr Leu Thr Arg Gly
 65 70 75 80
 Leu Lys His Leu Gly Pro Ser Phe His Val Leu Asp Ala Asn Arg Pro
 85 90 95
 Trp Leu Cys Tyr Trp Ile Ile His Ala Leu Ala Leu Leu Asp Glu Ile
 100 105 110
 Pro Asp Asp Val Glu Asp Asp Ile Val Asp Phe Leu Ser Arg Cys Gln
 115 120 125
 Asp Lys Asp Gly Gly Tyr Gly Gly Gly Pro Gly Gln Leu Pro His Leu
 130 135 140
 Ala Thr Thr Tyr Ala Ala Val Asn Thr Leu Val Thr Ile Gly Ser Glu
 145 150 155 160
 Arg Ala Leu Ser Ser Val Asn Arg Asp Asn Leu Tyr Lys Phe Met Leu
 165 170 175
 Arg Met Lys Asp Thr Ser Gly Ala Phe Arg Met His Asp Gly Gly Glu
 180 185 190
 Ile Asp Val Arg Ala Ser Tyr Thr Ala Ile Ser Val Ala Ser Leu Val
 195 200 205
 Asn Ile Leu Asp Gly Glu Leu Ala Lys Gly Val Gly Asn Tyr Ile Thr
 210 215 220
 Arg Cys Gln Thr Tyr Glu Gly Gly Ile Ala Gly Glu Pro Tyr Ala Glu
 225 230 235 240

Ala His Gly Gly Tyr Thr Phe Cys Gly Leu Ala Thr Met Ile Leu Leu
245 250 255

Asn Glu Val Asp Lys Leu Asp Leu Ala Ser Leu Ile Gly Trp Val Ala
260 265 270

Phe Arg Gln Gly Val Glu Cys Gly Phe Gln Gly Arg Thr Asn Lys Leu
275 280 285

Val Asp Gly Cys Tyr Ser Phe Trp Gln Gly Ala Ala Leu Ala Leu Thr
290 295 300

Val His Arg Val Ala Pro Thr Ala Lys
305 310

<210> 15
<211> 1504
<212> DNA
<213> Glycine max

<400> 15
gcacgaggac aaatcgcgcg ccgcgcgcgc cgtgtccgac ggtgagtaa cgtgagcaat 60
ggatggtaga gtcgcagggt ttccagattt accaaactctt cgccaccatt cctcgcaacg 120
cccaaaccct catgttggag ctccaacgcg ataatacatat gcagtagtgc tccaaaggcc 180
ttcgccatct cagttccgca tttccggttt tggacgctaa tcgaccctgg ctctgtctact 240
ggatcttcca ctccattgct ttgtcgggag aatccgtcga tgatgaactc gaagataacg 300
ctatcgattt tcttaaccgt tgcaggatc cgaatgggtg atagccggg ggaccaggcc 360
agatgctcta tattgccaca acttatgctg ctgttaattc acttattact ttgggtgggtg 420
agaaatccct ggcatacaatt aatagagata aactgtatgg gtttctcgcg cgagatgaagc 480
aaccaaatgg tggattcagg atgcatgatg aaggtgaaat tgatgttcga gcttgctaca 540
ctgccatttc tgttgcaagt gttttgaaca ttttggatga tgagctgac cagaatgttg 600
gagactacat tataagctgt caaacatatg aggttggcat tgcgtggtgag cctggttctg 660
aggctcatgg tgggtacacc ttttgggtat tagctacaat gattctgatt ggtgagggtta 720
atcacttgga tgcctctcga ttagtgtact ggggtggtatt ccgacaaggt aaggaatgtg 780
gattccaggg gagaacaaat aaactgtgtg atggatgcta ttccttttgg caggggagggtg 840
ctgttgctct attgcaaaaga ttatcttcta ttatcaacaa acagatggaa gagacatcac 900
agatttttgc ggtatcttat gtatctgaag caaaagaaag tttggatgga acctctagtc 960
atgcaacatg ccgtggtgag catgaaggca ccagtgaatc cagttcattc gatttttaaaa 1020
atattgccta taaatttatt aatgagtggg gagcacaaga accacttttt cacagtattg 1080
ctttacagca atatatcttc ttatgtgcac aggagcaaga ggtgtgactg agagacaaac 1140
cgggtaaaac tagagatcat tatcacacat gttactgttt aagtggactc tcattgtgccc 1200
agtatagttg gtcaaaagcac ccagattctc caaccactgcc taactctagta ttaggccccc 1260
actctaactc cttagaaccac atccaccccc tctttaatgt tgtcttgagg cgatatcgtg 1320
aagctcatga attctctctt actgagtcgt gaccactggt tttagctacc aacaacttta 1380
tttgtataat gtaaaataaa ttcatgtgaa catataaatg taaaacagca ttggattaaa 1440
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1500
aaaa 1504

<210> 16
<211> 429
<212> PRT
<213> Glycine max

<400> 16
Met Val Glu Ser Gln Val Phe Gln Ile Tyr Gln Leu Phe Ala Thr Ile
1 5 10 15
Pro Arg Asn Ala Gln Thr Leu Met Leu Glu Leu Gln Arg Asp Asn His
20 25 30

Met Gln Tyr Val Ser Lys Gly Leu Arg His Leu Ser Ser Ala Phe Ser
 35 40 45
 Val Leu Asp Ala Asn Arg Pro Trp Leu Cys Tyr Trp Ile Phe His Ser
 50 55 60
 Ile Ala Leu Ser Gly Glu Ser Val Asp Asp Glu Leu Glu Asp Asn Ala
 65 70 75 80
 Ile Asp Phe Leu Asn Arg Cys Gln Asp Pro Asn Gly Gly Tyr Ala Gly
 85 90 95
 Gly Pro Gly Gln Met Pro His Ile Ala Thr Thr Tyr Ala Ala Val Asn
 100 105 110
 Ser Leu Ile Thr Leu Gly Gly Glu Lys Ser Leu Ala Ser Ile Asn Arg
 115 120 125
 Asp Lys Leu Tyr Gly Phe Leu Arg Arg Met Lys Gln Pro Asn Gly Gly
 130 135 140
 Phe Arg Met His Asp Glu Gly Glu Ile Asp Val Arg Ala Cys Tyr Thr
 145 150 155 160
 Ala Ile Ser Val Ala Ser Val Leu Asn Ile Leu Asp Asp Glu Leu Ile
 165 170 175
 Gln Asn Val Gly Asp Tyr Ile Ile Ser Cys Gln Thr Tyr Glu Gly Gly
 180 185 190
 Ile Ala Gly Glu Pro Gly Ser Glu Ala His Gly Gly Tyr Thr Phe Cys
 195 200 205
 Gly Leu Ala Thr Met Ile Leu Ile Gly Glu Val Asn His Leu Asp Leu
 210 215 220
 Pro Arg Leu Val Asp Trp Val Val Phe Arg Gln Gly Lys Glu Cys Gly
 225 230 235 240
 Phe Gln Gly Arg Thr Asn Lys Leu Val Asp Gly Cys Tyr Ser Phe Trp
 245 250 255
 Gln Gly Gly Ala Val Ala Leu Leu Gln Arg Leu Ser Ser Ile Ile Asn
 260 265 270
 Lys Gln Met Glu Glu Thr Ser Gln Ile Phe Ala Val Ser Tyr Val Ser
 275 280 285
 Glu Ala Lys Glu Ser Leu Asp Gly Thr Ser Ser His Ala Thr Cys Arg
 290 295 300
 Gly Glu His Glu Gly Thr Ser Glu Ser Ser Ser Ser Asp Phe Lys Asn
 305 310 315 320
 Ile Ala Tyr Lys Phe Ile Asn Glu Trp Arg Ala Gln Glu Pro Leu Phe
 325 330 335
 His Ser Ile Ala Leu Gln Gln Tyr Ile Leu Leu Cys Ala Gln Glu Gln
 340 345 350

Glu Gly Gly Leu Arg Asp Lys Pro Gly Lys Arg Arg Asp His Tyr His
355 360 365

Thr Cys Tyr Cys Leu Ser Gly Leu Ser Leu Cys Gln Tyr Ser Trp Ser
370 375 380

Lys His Pro Asp Ser Pro Pro Leu Pro Asn Leu Val Leu Gly Pro Tyr
385 390 395 400

Ser Asn Leu Leu Glu Pro Ile His Pro Leu Phe Asn Val Val Leu Gly
405 410 415

Arg Tyr Arg Glu Ala His Glu Phe Phe Phe Thr Glu Ser
420 425

<210> 17
<211> 533
<212> DNA
<213> Glycine max

<220>
<221> unsure
<222> (499)

<220>
<221> unsure
<222> (525)

<400> 17
gagagagata cgaatccggc ggcggcgcca ccgtgtccga cggtagtca acgggaccag 60
tgatggttag agtcgcaggt gtttcagatt taccactct ttgccaccat tccctggcagc 120
gccccaaacc tcattgttaga gctgcaacgc gataatcaca tgcagtatct ctccaaaggc 180
ctacgccatc tcagttccgc gttttctgtc ttggacgcta atcgaccctg gctctgttac 240
tgatctttcc attccattgc tttgctggga gaatccgtcg acgacgaact cgaagataac 300
actatcgatt ttcttaaccg ttgccaggat ccgaatggtg gatatgctgg gggaccaggc 360
cagatgcctc acattgccac aacatatgct gcagttaata cacttattac tttgggtggt 420
cagaaatcct ggcatacaat aataggtgag ataaactgta tgggtttctg cgccggatga 480
agcaatcaaa tggggggant caagatgcac gatgaaagga gaaanttgat gtc 533

<210> 18
<211> 141
<212> PRT
<213> Glycine max

<400> 18
Asp Thr Asn Pro Ala Ala Pro Pro Cys Pro Thr Val Ser Gln Arg
1 5 10 15

Asp Gln Trp Met Val Glu Ser Gln Val Phe Gln Ile Tyr Gln Leu Phe
20 25 30

Ala Thr Ile Pro Gly Ser Ala Gln Asn Leu Met Leu Glu Leu Gln Arg
35 40 45

Asp Asn His Met Gln Tyr Leu Ser Lys Gly Leu Arg His Leu Ser Ser
50 55 60

Ala Phe Ser Val Leu Asp Ala Asn Arg Pro Trp Leu Cys Tyr Trp Ile
65 70 75 80

Phe His Ser Ile Ala Leu Leu Gly Glu Ser Val Asp Asp Glu Leu Glu
85 90 95

Asp Asn Thr Ile Asp Phe Leu Asn Arg Cys Gln Asp Pro Asn Gly Gly
100 105 110

Tyr Ala Gly Gly Pro Gly Gln Met Pro His Ile Ala Thr Thr Tyr Ala
115 120 125

Ala Val Asn Thr Leu Ile Thr Leu Gly Gly Gln Lys Ser
130 135 140

<210> 19

<211> 333

<212> PRT

<213> Pisum sativum

<400> 19

Met Ala Gly Asn Ile Glu Val Glu Glu Asp Asp Arg Val Pro Leu Arg
1 5 10 15

Leu Arg Pro Glu Trp Ser Asp Val Thr Pro Ile Pro Gln Asp Asp Gly
20 25 30

Pro Ser Pro Val Val Pro Ile Asn Tyr Ser Glu Glu Phe Ser Glu Val
35 40 45

Met Asp Tyr Phe Arg Ala Val Tyr Phe Ala Lys Glu Leu Ser Ser Arg
50 55 60

Ala Leu Ala Leu Thr Ala Glu Ala Ile Gly Leu Asn Ala Gly Asn Tyr
65 70 75 80

Thr Val Trp His Phe Arg Arg Leu Leu Leu Glu Ser Leu Lys Val Asp
85 90 95

Leu His Val Glu Arg Glu Phe Val Glu Arg Val Ala Ser Gly Asn Ser
100 105 110

Lys Asn Tyr Gln Ile Trp His His Arg Arg Trp Val Ala Glu Lys Leu
115 120 125

Gly Pro Glu Ala Arg Asn Ser Glu Leu Glu Phe Thr Lys Lys Ile Leu
130 135 140

Ser Val Asp Ala Lys His Tyr His Ala Trp Ser His Arg Gln Trp Val
145 150 155 160

Leu Gln Asn Leu Gly Gly Trp Glu Asp Glu Leu Ser Tyr Cys Ser Glu
165 170 175

Leu Leu Ala Glu Asp Ile Phe Asn Asn Ser Ala Trp Asn Gln Arg Tyr
180 185 190

Phe Val Ile Thr Arg Ser Pro Val Leu Gly Gly Leu Lys Ala Met Arg
195 200 205

Glu Ser Glu Val Leu Phe Thr Val Glu Ala Ile Ile Ser Tyr Pro Glu
210 215 220

Asn Glu Ser Ser Trp Arg Tyr Leu Arg Gly Leu Phe Lys Asp Glu Ser
 225 230 235 240
 Thr Leu Tyr Val Asn Asp Ala Gln Val Ser Ser Leu Cys Leu Lys Ile
 245 250 255
 Leu Lys Thr Lys Ser Asn Tyr Leu Phe Ala Leu Ser Thr Leu Leu Asp
 260 265 270
 Leu Ser Ala Ser Val Ile Gln Pro Asn Glu Asp Phe Arg Asp Ala Ile
 275 280 285
 Glu Ala Leu Arg Leu Gln Ile Leu Ile Lys Gln Asp Ser Asp Ile Ala
 290 295 300
 Ile Thr Ile Cys Ser Ile Leu Glu Gln Val Asp Pro Ile Arg Val Asn
 305 310 315 320
 Tyr Trp Val Trp Arg Lys Ser Arg Leu Pro Gln Ala Ala
 325 330

<210> 20
 <211> 326
 <212> PRT
 <213> Arabidopsis thaliana

<400> 20
 Met Asn Phe Asp Glu Thr Val Pro Leu Ser Gln Arg Leu Glu Trp Ser
 1 5 10 15
 Asp Val Val Pro Leu Thr Gln Asp Asp Gly Pro Asn Pro Val Val Pro
 20 25 30
 Ile Ala Tyr Lys Glu Glu Phe Arg Glu Thr Met Asp Tyr Phe Arg Ala
 35 40 45
 Ile Tyr Phe Ser Asp Glu Arg Ser Pro Arg Ala Leu Arg Leu Thr Glu
 50 55 60
 Glu Thr Leu Leu Leu Asn Ser Gly Asn Tyr Thr Val Trp His Phe Arg
 65 70 75 80
 Arg Leu Val Leu Glu Ala Leu Asn His Asp Leu Phe Glu Glu Leu Glu
 85 90 95
 Phe Ile Glu Arg Ile Ala Glu Asp Asn Ser Lys Asn Tyr Gln Leu Trp
 100 105 110
 His His Arg Arg Trp Val Ala Glu Lys Leu Gly Pro Asp Val Ala Gly
 115 120 125
 Arg Glu Leu Glu Phe Thr Arg Arg Val Leu Ser Leu Asp Ala Lys His
 130 135 140
 Tyr His Ala Trp Ser His Arg Gln Trp Thr Leu Arg Ala Leu Gly Gly
 145 150 155 160
 Trp Glu Asp Glu Leu Asp Tyr Cys His Glu Leu Leu Glu Ala Asp Val
 165 170 175

Phe Asn Asn Ser Ala Trp Asn Gln Arg Tyr Tyr Val Ile Thr Gln Ser
 180 185 190
 Pro Leu Leu Gly Gly Leu Glu Ala Met Arg Glu Ser Glu Val Ser Tyr
 195 200 205
 Thr Ile Lys Ala Ile Leu Thr Asn Pro Ala Asn Glu Ser Ser Trp Arg
 210 215 220
 Tyr Leu Lys Ala Leu Tyr Lys Asp Asp Lys Glu Ser Trp Ile Ser Asp
 225 230 235 240
 Pro Ser Val Ser Ser Val Cys Leu Asn Val Leu Ser Arg Thr Asp Cys
 245 250 255
 Phe His Gly Phe Ala Leu Ser Thr Leu Leu Asp Leu Leu Cys Asp Gly
 260 265 270
 Leu Arg Pro Thr Asn Glu His Lys Asp Ser Val Arg Ala Leu Ala Asn
 275 280 285
 Glu Glu Pro Glu Thr Asn Leu Ala Asn Leu Val Cys Thr Ile Leu Gly
 290 295 300
 Arg Val Asp Pro Ile Arg Ala Asn Tyr Trp Ala Trp Arg Lys Ser Lys
 305 310 315 320
 Ile Thr Val Ala Ala Ile
 325
 <210> 21
 <211> 470
 <212> PRT
 <213> Lycopersicon esculentum
 <400> 21
 Met Glu Ser Arg Lys Val Thr Lys Thr Leu Glu Asp Gln Trp Val Val
 1 5 10 15
 Glu Arg Arg Val Arg Glu Ile Tyr Asp Tyr Phe Tyr Ser Ile Ser Pro
 20 25 30
 Asn Ser Pro Ser Asp Leu Ile Glu Ile Glu Arg Asp Lys His Phe Gly
 35 40 45
 Tyr Leu Ser Gln Gly Leu Arg Lys Leu Gly Pro Ser Phe Ser Val Leu
 50 55 60
 Asp Ala Ser Arg Pro Trp Leu Cys Tyr Trp Thr Leu His Ser Ile Ala
 65 70 75 80
 Leu Leu Gly Glu Ser Ile Gly Gly Lys Leu Glu Asn Asp Ala Ile Asp
 85 90 95
 Phe Leu Thr Arg Cys Gln Asp Lys Asp Gly Gly Tyr Gly Gly Gly Pro
 100 105 110
 Gly Gln Met Pro His Leu Ala Thr Thr Tyr Ala Ala Val Asn Ser Leu
 115 120 125

Ile Thr Leu Gly Lys Pro Glu Ala Leu Ser Ser Ile Asn Arg Glu Lys
 130 135 140
 Leu Tyr Thr Phe Leu Leu Arg Met Lys Asp Ala Ser Gly Gly Phe Arg
 145 150 155 160
 Met His Asp Gly Gly Glu Val Asp Val Arg Ala Cys Tyr Thr Ala Ile
 165 170 175
 Ser Val Ala Asn Ile Leu Asn Ile Val Asp Asp Glu Leu Ile His Gly
 180 185 190
 Val Gly Asn Tyr Ile Leu Ser Cys Gln Thr Tyr Glu Gly Gly Ile Ala
 195 200 205
 Gly Glu Pro Gly Ser Glu Ala His Gly Gly Tyr Thr Phe Cys Gly Leu
 210 215 220
 Ala Ala Met Ile Leu Ile Asn Glu Val Asp Arg Leu Asp Leu Pro Gly
 225 230 235 240
 Leu Ile Asp Trp Val Val Phe Arg Gln Gly Val Glu Gly Gly Phe Gln
 245 250 255
 Gly Arg Thr Asn Lys Leu Val Asp Gly Cys Tyr Ser Phe Trp Gln Gly
 260 265 270
 Ala Val Val Phe Leu Ile Gln Arg Leu Asn Leu Ile Val His Glu Gln
 275 280 285
 Leu Gly Leu Ser Asn Asp Leu Ser Thr Glu Ser Ala Asp Asp Ser Ser
 290 295 300
 Glu Ser Glu Leu Ser Asp Glu Glu Glu His Leu Glu Gly Ile Ser Ser
 305 310 315 320
 His Val Gln Asp Thr Phe Pro Leu Gly Gln Ala Gly Ala Cys Gln Glu
 325 330 335
 Asn Ala Ser His Ser Pro Lys Ile Ala Asp Thr Gly Tyr Glu Phe Ile
 340 345 350
 Asn Arg Pro Ile Ala Met Arg Pro Leu Phe Asp Ser Met Tyr Leu Gln
 355 360 365
 Gln Tyr Val Leu Leu Cys Ser Gln Ile Glu Val Gly Gly Phe Arg Asp
 370 375 380
 Lys Pro Gly Lys Gly Arg Asp Tyr Tyr His Thr Cys Tyr Cys Leu Ser
 385 390 395 400
 Gly Leu Ser Ile Ala Gln Tyr Ser Trp Thr Asp Glu Ala Asp Ser Thr
 405 410 415
 Pro Leu Pro Arg Asp Val Phe Gly Pro Tyr Ser Lys Cys Leu Leu Glu
 420 425 430
 Gln Val His Pro Leu Phe Asn Val Val Leu Asp Arg Tyr Tyr Glu Ala
 435 440 445

Arg Glu Tyr Ser Gln Ala Cys Glu Thr Val Ser Pro Leu Ser Leu Ala
 450 455 460

Pro Thr Phe Ser Glu Thr
 465 470

<210> 22
 <211> 419
 <212> PRT
 <213> Pisum sativum

<400> 22
 Met Glu Ala Ser Thr Ala Ala Glu Thr Pro Thr Pro Thr Val Ser Gln
 1 5 10 15

Arg Asp Gln Trp Ile Val Glu Ser Gln Val Phe His Ile Tyr Gln Leu
 20 25 30

Phe Ala Asn Ile Pro Pro Asn Ala Gln Ser Ile Ile Arg Pro Trp Leu
 35 40 45

Cys Tyr Trp Ile Ile His Ser Ile Ala Leu Leu Gly Glu Ser Ile Asp
 50 55 60

Asp Asp Leu Glu Asp Asn Thr Val Asp Phe Leu Asn Arg Cys Gln Asp
 65 70 75 80

Pro Asn Gly Gly Tyr Ala Gly Gly Pro Gly Gln Met Pro His Leu Ala
 85 90 95

Thr Thr Tyr Ala Ala Val Asn Thr Leu Ile Thr Leu Gly Gly Glu Lys
 100 105 110

Ser Leu Ala Ser Ile Asn Arg Asn Lys Leu Tyr Gly Phe Met Arg Arg
 115 120 125

Met Lys Gln Pro Asn Gly Gly Phe Arg Met His Asp Glu Gly Glu Ile
 130 135 140

Asp Val Arg Ala Cys Tyr Thr Ala Ile Ser Val Ala Ser Val Leu Asn
 145 150 155 160

Ile Leu Asp Asp Glu Leu Ile Lys Asn Val Gly Asp Phe Ile Leu Ser
 165 170 175

Cys Gln Thr Tyr Glu Gly Gly Leu Ala Gly Glu Pro Gly Ser Glu Ala
 180 185 190

His Gly Gly Tyr Thr Phe Cys Gly Leu Ala Ala Met Ile Leu Ile Gly
 195 200 205

Glu Val Asn Arg Leu Asp Leu Pro Arg Leu Leu Asp Trp Val Val Phe
 210 215 220

Arg Gln Gly Lys Glu Cys Gly Phe Gln Gly Arg Thr Asn Lys Leu Val
 225 230 235 240

Asp Gly Cys Tyr Ser Phe Trp Gln Gly Gly Ala Val Ala Leu Leu Gln
 245 250 255

Arg Leu His Ser Ile Ile Asp Glu Gln Met Ala Glu Ala Ser Gln Phe
 260 265 270

Val Thr Val Ser Asp Ala Pro Glu Glu Lys Glu Cys Leu Asp Gly Thr
 275 280 285

Ser Ser His Ala Thr Ser His Ile Arg His Glu Gly Met Asn Glu Ser
 290 295 300

Cys Ser Ser Asp Val Lys Asn Ile Gly Tyr Asn Phe Ile Ser Glu Trp
 305 310 315 320

Arg Gln Ser Glu Pro Leu Phe His Ser Ile Ala Leu Gln Gln Tyr Ile
 325 330 335

Leu Leu Cys Ser Gln Glu Gln Asp Gly Gly Leu Arg Asp Lys Pro Gly
 340 345 350

Lys Arg Arg Asp His Tyr His Ser Cys Tyr Cys Leu Ser Gly Leu Ser
 355 360 365

Leu Cys Gln Tyr Ser Trp Ser Lys Arg Pro Asp Ser Pro Pro Leu Pro
 370 375 380

Lys Val Val Met Gly Pro Tyr Ser Asn Leu Leu Glu Pro Ile His Pro
 385 390 395 400

Leu Phe Asn Val Val Leu Asp Arg Tyr Arg Glu Ala His Glu Phe Phe
 405 410 415

Ser Gln Leu

<210> 23
 <211> 419
 <212> PRT
 <213> Pisum sativum

<400> 23
 Met Glu Ala Ser Thr Ala Ala Glu Thr Pro Thr Pro Thr Val Ser Gln
 1 5 10 15

Arg Asp Gln Trp Ile Val Glu Ser Gln Val Phe His Ile Tyr Gln Leu
 20 25 30

Phe Ala Asn Ile Pro Pro Asn Ala Gln Ser Ile Ile Arg Pro Trp Leu
 35 40 45

Cys Tyr Trp Ile Ile His Ser Ile Ala Leu Leu Gly Glu Ser Ile Asp
 50 55 60

Asp Asp Leu Glu Asp Asn Thr Val Asp Phe Leu Asn Arg Cys Gln Asp
 65 70 75 80

Pro Asn Gly Gly Tyr Ala Gly Gly Pro Gly Gln Met Pro His Leu Ala
 85 90 95

Thr Thr Tyr Ala Ala Val Asn Thr Leu Ile Thr Leu Gly Gly Glu Lys
 100 105 110

Ser Leu Ala Ser Ile Asn Arg Asn Lys Leu Tyr Gly Phe Met Arg Arg
 115 120 125
 Met Lys Gln Pro Asn Gly Gly Phe Arg Met His Asp Glu Gly Glu Ile
 130 135 140
 Asp Val Arg Ala Cys Tyr Thr Ala Ile Ser Val Ala Ser Val Leu Asn
 145 150 155 160
 Ile Leu Asp Asp Glu Leu Ile Lys Asn Val Gly Asp Phe Ile Leu Ser
 165 170 175
 Cys Gln Thr Tyr Glu Gly Gly Leu Ala Gly Glu Pro Gly Ser Glu Ala
 180 185 190
 His Gly Gly Tyr Thr Phe Cys Gly Leu Ala Ala Met Ile Leu Ile Gly
 195 200 205
 Glu Val Asn Arg Leu Asp Leu Pro Arg Leu Leu Asp Trp Val Val Phe
 210 215 220
 Arg Gln Gly Lys Glu Cys Gly Phe Gln Gly Arg Thr Asn Lys Leu Val
 225 230 235 240
 Asp Gly Cys Tyr Ser Phe Trp Gln Gly Gly Ala Val Ala Leu Leu Gln
 245 250 255
 Arg Leu His Ser Ile Ile Asp Glu Gln Met Ala Glu Ala Ser Gln Phe
 260 265 270
 Val Thr Val Ser Asp Ala Pro Glu Glu Lys Glu Cys Leu Asp Gly Thr
 275 280 285
 Ser Ser His Ala Thr Ser His Ile Arg His Glu Gly Met Asn Glu Ser
 290 295 300
 Cys Ser Ser Asp Val Lys Asn Ile Gly Tyr Asn Phe Ile Ser Glu Trp
 305 310 315 320
 Arg Gln Ser Glu Pro Leu Phe His Ser Ile Ala Leu Gln Gln Tyr Ile
 325 330 335
 Leu Leu Cys Ser Gln Glu Gln Asp Gly Gly Leu Arg Asp Lys Pro Gly
 340 345 350
 Lys Arg Arg Asp His Tyr His Ser Cys Tyr Cys Leu Ser Gly Leu Ser
 355 360 365
 Leu Cys Gln Tyr Ser Trp Ser Lys Arg Pro Asp Ser Pro Pro Leu Pro
 370 375 380
 Lys Val Val Met Gly Pro Tyr Ser Asn Leu Leu Glu Pro Ile His Pro
 385 390 395 400
 Leu Phe Asn Val Val Leu Asp Arg Tyr Arg Glu Ala His Glu Phe Phe
 405 410 415
 Ser Gln Leu